

60th Annual Scientific Session & Expo

E1039

JACC April 5, 2011

Volume 57, Issue 14



MYOCARDIAL ISCHEMIA AND INFARCTION

EFFECT OF APIXABAN, AN ORAL, DIRECT AND SELECTIVE FACTOR XA INHIBITOR ON INFLAMMATORY BIOMARKERS FOLLOWING ACUTE CORONARY SYNDROME

ACC Poster Contributions

Ernest N. Morial Convention Center, Hall F

Monday, April 04, 2011, 9:30 a.m.-10:45 a.m.

Session Title: Management of ACS

Abstract Category: 4. Unstable Ischemic Syndrome/Long-Term Outcome

Session-Poster Board Number: 1073-366

Authors: *Richard C. Becker, Annie Lin, Hongqiu Yang, Yuchen Barrett, Puneet Mohan, Jessie Wang, John Alexander, Duke Clinical Research Institute, Durham, NC, Bristol-Myers Squibb, Princeton, NJ*

Background: Apixaban is an oral, direct and selective factor Xa inhibitor under development for secondary prevention in acute coronary syndrome (ACS).

Methods: Apixaban's effect on biomarkers of inflammation and coagulant regulatory proteins was assessed in a randomized, double-blinded, placebo-controlled phase 2 dose-ranging trial. Samples were obtained at baseline, week 3 and week 26 in 1715 patients with ST-segment elevation or non-ST elevation ACS who received apixaban (5-20mg daily) or placebo. Apixaban plasma concentrations were measured directly by LC/MS/MS, and anti-Xa activity was determined with a chromogenic assay using an apixaban calibrator.

Results: There was a strong, direct and linear relationship between apixaban plasma concentration and anti-Xa activity ($r = 0.97$). High sensitivity (hs) C-reactive protein (CRP) ($p < 0.0001$) fibrinogen, ($p < 0.0001$), soluble CD-40Ligand (sCD40L) ($p < 0.0001$) and myeloperoxidase (MPO) ($p = 0.00001$) decreased over time from elevated baseline values in all groups. Tissue factor pathway inhibitor (TFPI) increased over time ($p < 0.0001$). The changes were greater at week 26 compared with week 3 across increasing quartiles of apixaban plasma concentration and anti-Xa activity (figure). An apixaban level between 100 and 200ng/ml, achieved with the 10mg total daily doses investigated in APPRAISE-2, AVEROES, and ARISTOTLE was associated with maximal suppression of hs-CRP ($p = 0.03$), CD40L ($p = 0.01$) and MPO ($p = 0.04$) at 26 weeks of treatment.

Conclusions: Apixaban - an oral, direct and selective factor Xa inhibitor, in addition to its predicted anticoagulant effects may also attenuate leukocyte- and platelet-mediated inflammation and upregulate intrinsic anticoagulant activity, supporting the pleiotropic properties of factor Xa.